



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

(*Phlæotomus pileatus abieticola*) in this State is May 10-15, consequently I was surprised to find a nest on May 20, 1918, in northern Huntingdon County containing three nestlings about one week old. The nest was seventy feet up in the dead top of a large rock oak in a thick forest.

Judging by the age of the young birds and allowing eighteen days for the incubation of the eggs, this early pair must have had a fresh set about April 25.

Three other nests found by Richard C. Harlow and the writer this year, in the same county held fresh and slightly incubated eggs on May 16 and 17, respectively.—RICHARD F. MILLER, *Philadelphia, Pa.*

Relative Length of the Intestinal Cæca in Trogons.—In his article on 'The Anatomy of the Cuban Trogon' in 'The Auk' for July, 1918 (p. 286), Dr. H. L. Clark records the length of the intestines and cæca of this trogon and remarks "The cæca are thus relatively very long, much longer than in the species of Trogon and *Pharomacrus* examined by Garrod."

I have examined the viscera of twelve specimens, representing six species, of Central American Trogons. These with the four individuals recorded by Garrod and Clark give us for comparison a total of eight species belonging to three very distinct groups of Neotropical Trogonidæ. The following figures express the ratio of the length of the cæca to that of the intestines, the latter being represented by 100.

<i>Pharomachrus mocinno</i>	(7 specimens),	9.3
<i>Trogonurus mexicanus</i>	(1 specimen),	10.6
" <i>puella</i>	(2 specimens),	13.8
" <i>curucui</i>	(1 specimen),	14.6
<i>Chrysotrogon caligatus</i>	(1 specimen),	16.2
<i>Trogonurus elegans</i>	(2 specimens),	16.7
<i>Trogon melanocephalus</i>	(1 specimen),	17.2
<i>Priotelus temnurus</i>	(1 specimen),	17.8

Thus in *Pharomachrus* the cæca average slightly more than one-eleventh of the total length of the intestinal tract, while in *Priotelus* they exceed one-sixth of the intestinal length. In *Pharomachrus* they are relatively shorter than in the other genera but *Trogonurus mexicanus* connects the two groups. The figures indicate that the cæca of *Priotelus* are a trifle longer than those of *Trogonurus*, *Chrysotrogon* and *Trogon*, but there is great individual variation in the length of these appendages and additional specimens will undoubtedly show that there is at most only a slight average difference.—W. DEW. MILLER, *American Museum of Natural History, New York City.*

The Range and Status of *Aphelocoma californica hypoleuca* Ridgway.

—As information supplemental to Mr. H. S. Swarth's excellent revision